

Sunlight


PHYS 4400, Principles and Varieties of Solar Energy

**Instructor: Randy J. Ellingson
The University of Toledo**


January 22, 2015



Solar Radiation Spectrum



Earth's Motion around the Sun



δ = angle of declination between Sun's rays and the plane of the equator

From "Solar Cells", by C. Hu and R.M. White



After Coordinate Transformation (earth-centric view)




Figure 2.1 (a) The conventional sun-centered view of the sun-earth system. (b) An earth-centered view, which is easier to visualize. For example, the declination angle δ between the sun ray and the plane of the equator is better illustrated in *b*. The date given may vary by one day or so.

From "Solar Cells", by C. Hu and R.M. White

Daily Variation in Energy Collection





Figure 2.7 Qualitative plots of seasonal variations of daily solar energy collection for a location north of 23.5°N. "Tilt" (see Sec. 2.5) is the angle between the collector plate and the horizontal plane. The plate is tilted due south.

From "Solar Cells", by C. Hu and R.M. White





Solar Constant

We've been informed that the total luminous power output from the Sun is 3.846×10^{26} W.

Now we'd like to determine the portion of that power incident on Earth, and the average areal intensity (i.e., averaged over the area of Earth's surface).

From in-class exercises, we figured out the solar constant ($\sim 1.75 \times 10^{17}$ W).

We found that the intensity of sunlight arriving at the top of Earth's atmosphere is 1.375 kW m^{-2} .

We determined the intensity of arriving sunlight averaged over Earth's total surface area to be 343 W m^{-2} . (averaged over the total area of the Earth, and averaged over day and night).

$$\begin{aligned} r_{\text{Earth}} &= 6.37 \times 10^6 \text{ m} \\ 1 \text{ AU} &= 1.496 \times 10^{11} \text{ m} \end{aligned}$$



Standard solar spectra

Sunlight varies with factors including location, weather, and the time of day.

Standard solar spectra are defined to enable comparison of materials and device performance among manufacturers and laboratories around the world.

There are standard for extraterrestrial and terrestrial sunlight:

Extraterrestrial: this spectrum, referred to as AM0, defines sunlight just above Earth's atmosphere, where the sunlight has an intensity dependent on the solar constant.


Terrestrial: refers to sunlight which has passed through Earth's atmosphere. There are two standard solar spectra, referred to as AM1.5G (G stands for Global Tilt), and AM1.5 Direct (+circumsolar).







AM0: the spectrum above Earth's atmosphere

The AM0 spectrum applies to satellites and high-flying aircraft, which access the spectrum prior to any influence from Earth's atmosphere.

Integrated spectral irradiance = 1366 W/m².




Atmospheric effects on sunlight



Air Mass

$$AM = \frac{1}{\cos \theta}$$




Does $AM = \infty$ when $\theta = 90^\circ$?

$$AM = \frac{1}{\cos \theta + 0.5057(96.07995 - \theta)^{-1.6364}}$$



AM1.5G: reference spectrum including direct and diffuse sunlight

- AM 1.5: From the equation provided for Air Mass, one calculates that $\cos \theta = 0.667$, so that $\theta = 48.2^\circ$. This represents the zenith angle, that is, the angle relative to the direction normal to Earth's surface.
- Toledo latitude: 41.6639° N
- Integrating the energy within AM1.5G yields 1000 W m^{-2} .



AM1.5D: reference spectrum including direct and circumsolar diffuse sunlight

Includes sunlight scattered within 2.5° angle of the direct sunlight. Used for clear days, relevant to concentrated sunlight applications for which the ability to focus the sunlight depends critically on the direction the light is traveling.

$$I_G = 1.1 \cdot I_D$$

