


# X-ray Diffraction and Crystal Structures

Week of March 22, 2010

**Modern Physics Laboratory**  
**(Physics 6180/7180)**


The University of Toledo  
Instructor: Randy Ellingson



## X-Ray Generation

X-rays are electromagnetic radiation with wavelength  $\sim 1 \text{ \AA} = 10^{-10} \text{ m}$   
(visible light  $\sim 5.5 \times 10^{-7} \text{ m}$ )

X-ray generation



X-ray wavelengths too short to be resolved by a standard optical grating

$$\theta = \sin^{-1} \frac{m\lambda}{d} = \sin^{-1} \frac{(1)(0.1 \text{ nm})}{3000 \text{ nm}} = 0.0019^\circ$$


## X-Ray Generation

The most common metal used is copper, which can be kept cool easily, due to its high thermal conductivity, and which produces strong  $K_\alpha$  and  $K_\beta$  lines. The  $K_\beta$  line is sometimes suppressed with a thin ( $\sim 10\ \mu\text{m}$ ) nickel foil.

- **K-alpha ( $K_\alpha$ )** emission lines result when an electron transitions to the innermost "K" shell (principal quantum number 1) from a 2p orbital of the second or "L" shell (with principal quantum number 2).
- The  $K_\alpha$  line is actually a doublet, with slightly different energies depending on spin-orbit interaction energy between the electron spin and the orbital momentum of the 2p orbital.


$$\lambda(K_\alpha) = 0.154\ \text{nm}$$

$$\lambda(K_\beta) = 0.139\ \text{nm}$$




Atomic levels involved in copper  $K_\alpha$  and  $K_\beta$  emission.


## $K_{\alpha}$ and $K_{\beta}$ X-ray lines



from Preston and Dietz, p. 191.

## X-Ray diffraction







## X-Ray Diffraction -- Bragg's Law


Diffraction of x-rays by crystal: spacing  $d$  of adjacent crystal planes on the order of 0.1 nm

→ three-dimensional diffraction grating with diffraction maxima along angles where reflections from different planes interfere constructively




$$2d \sin \theta = m\lambda \text{ for } m = 0, 1, 2, \dots$$

**Bragg's Law**



## The Braggs (Bragg's Law)



**Sir William Henry Bragg**  
**1862-1942**




**William Lawrence Bragg**  
**1890-1971**

Bragg occupied the Cavendish chair of physics at the University of Leeds from 1909. He continued his work on X-rays with much success. He invented the X-ray spectrometer and with his son, William Lawrence Bragg, then a research student at Cambridge, founded the new science of X-ray analysis of crystal structure.

In 1915 father and son were jointly awarded the Nobel Prize in Physics for their studies, using the X-ray spectrometer, of X-ray spectra, X-ray diffraction, and of crystal structure.

## X-Ray Diffraction, cont'd



Interplanar spacing  $d$  is related to the unit cell dimension  $a_0$



$$5d = \sqrt{\frac{5}{4}} a_0 \quad \text{or} \quad d = \frac{a_0}{20} = 0.2236a_0$$

Not only can crystals be used to separate different x-ray wavelengths, but x-rays in turn can be used to study crystals, for example determine the type of crystal ordering and  $a_0$ .




## Crystal structure, lattice planes, and Miller indices


Planes with different Miller indices in cubic crystals. The *inverse* of these fractional intercepts yields the Miller indices  $h, k, l$ .




## Crystal structure and Miller indices



(001)




(100)




(010)


Planes with different Miller indices in cubic crystals.




(101)




(110)




(011)



(111)




(1 $\bar{1}\bar{1}$ )



( $\bar{1}\bar{1}\bar{1}$ )


# Crystal structure and Miller indices



**Indexing lattice planes**

[http://www.msm.cam.ac.uk/doitpoms/tlplib/miller\\_indices/lattice\\_index.php](http://www.msm.cam.ac.uk/doitpoms/tlplib/miller_indices/lattice_index.php)

## Rock salt (cubic) crystal structure



$$d_{hkl} = \frac{a_0}{\sqrt{h^2 + k^2 + l^2}}$$

**Structure factor for NaCl:**



$$F = \left[ f_{Na} + f_{Cl} e^{i\pi(h+k+l)} \right] \left[ 1 + e^{i\pi(h+k)} + e^{i\pi(h+l)} + e^{i\pi(k+l)} \right]$$

$$F = 4(f_{Na} + f_{Cl}) \quad \text{if } h, k, l \text{ are even}$$

$$F = 4(f_{Na} - f_{Cl}) \quad \text{if } h, k, l \text{ are odd}$$


$$F = 0 \quad \text{if } h, k, l \text{ are mixed}$$

## X-Ray diffraction (XRD) pattern (diffractogram) from NaCl



$$d_{hkl} = \frac{a_0}{\sqrt{h^2 + k^2 + l^2}}$$

## LiF diffractogram (Cu $K_\alpha$ )



# TEL-X-Ometer



$K_{\alpha 1}$	1.540 Å
$K_{\alpha 2}$	1.544 Å
$K_{\beta}$	1.392 Å