

## **Answer Key**

1. D
2. E
3. E
4. B


MIDTERM #1

Solution:

5. The charges are equidistant from the point where we are evaluating the potential — which is computed using Eq. 24-27 (or its integral equivalent). Eq. 24-27 implicitly assumes  $V \rightarrow 0$  as  $r \rightarrow \infty$ . Thus, we have

$$V = \frac{1}{4\pi\epsilon_0} \frac{+Q_1}{R} + \frac{1}{4\pi\epsilon_0} \frac{-2Q_1}{R} + \frac{1}{4\pi\epsilon_0} \frac{+3Q_1}{R} = \frac{1}{4\pi\epsilon_0} \frac{2Q_1}{R} = \frac{2(8.99 \times 10^9)(4.52 \times 10^{-12})}{0.0850} = 0.956 \text{ V.}$$

6. We use a Gaussian surface in the form of a box with rectangular sides. The cross section is shown with dashed lines in the diagram below.



It is centered at the central plane of the slab, so the left and right faces are each a distance  $x$  from the central plane. We take the thickness of the rectangular solid to be  $a$ , the same as its length, so the left and right faces are squares. The electric field is normal to the left and right faces and is uniform over them. Since  $\rho = 5.80 \text{ fC/m}^3$  is positive, it points outward at both faces: toward the left at the left face and toward the right at the right face. Furthermore, the magnitude is the same at both faces. The electric flux through each of these faces is  $Ea^2$ . The field is parallel to the other faces of the Gaussian surface and the flux through them is zero. The total flux through the Gaussian surface is  $\Phi = 2Ea^2$ . The volume enclosed by the Gaussian surface is  $2a^2x$  and the charge contained within it is  $q = 2a^2x\rho$ . Gauss' law yields  $2\epsilon_0Ea^2 = 2a^2x\rho$ . We solve for the magnitude of the electric field:  $E = \rho x / \epsilon_0$ .

(a) For  $x=0$ ,  $E=0$ .

(b) For  $x = 2.00 \text{ mm} = 2.00 \times 10^{-3} \text{ m}$ ,

$$E = (5.80 \times 10^{-15})(2.00 \times 10^{-3}) / (8.85 \times 10^{-12}) = 1.31 \times 10^{-6} \text{ N/C}.$$

(c) For  $x = d/2 = 4.70 \text{ mm} = 4.70 \times 10^{-3} \text{ m}$ ,

$$E = (5.80 \times 10^{-15})(4.70 \times 10^{-3}) / (8.85 \times 10^{-12}) = 3.08 \times 10^{-6} \text{ N/C}.$$

(d) For  $x = 26.0 \text{ mm} = 2.60 \times 10^{-2} \text{ m}$ , we take a Gaussian surface of the same shape and orientation, but with  $x > d/2$ , so the left and right faces are outside the slab. The total flux through the surface is again  $\Phi = 2Ea^2$  but the charge enclosed is now  $q = a^2 d\rho$ . Gauss' law yields  $2\epsilon_0 E a^2 = a^2 d\rho$ , so

$$E = \frac{\rho d}{2\epsilon_0} = \frac{(5.80 \times 10^{-15})(9.40 \times 10^{-3})}{2(8.85 \times 10^{-12})} = 3.08 \times 10^{-6} \text{ N/C}.$$